

The Nuclear Bulge extinction

Andrew J. Gosling, Katherine M. Blundell

Oxford Astrophysics, Denys Wilkinson Building, Keble Road, Oxford, OX1 3RH

Reba M. Bandyopadhyay

*Department of Astronomy, University of Florida, 210 Bryant Space Science Center,
 Gainesville, FL 32611-2055*

Phil Lucas

School of Astronomy, University of Hertfordshire, College Lane, Hatfield, Herts AL10 9AB

Abstract. We present a new, high resolution ($5''$ per pixel) near-infrared extinction map of the Nuclear Bulge using data from the UKIDSS-GPS. Using photometry from the J , H and K -bands we show that the extinction law parameter α is also highly variable in this region on similar scales to the absolute extinction. We show that only when this extinction law variation is taken into account can the extinction be measured consistently at different wavelengths.

1. Introduction

Study of the Nuclear Bulge (NB) is extremely difficult as it is one of the most highly obscured regions of the Galaxy. The extinction towards the NB is so great that it is almost impossible to observe at visual wavelengths. It is necessary to observe in the near-infrared (NIR) to obtain imaging similar to that normally obtained in visual. Previous measures of the NIR extinction towards the NB (Catchpole et al. 1990; Schultheis et al. 1999; Dutra et al. 2003) found the extinction to be highly spatially variable. We present a new mapping of the extinction towards the NB at a much higher resolution than previously possible.

2. Extinction mapping

We mapped the extinction across a region of approximately $2^\circ \times 2^\circ$ centred on Sgr A* using near-infrared data from the (United Kingdom Infrared Deep Sky Survey - Galactic Plane Survey) UKIDSS-GPS (Lucas et al. 2007). Using a 2:1 over-sampling, we obtained a resolution of $5''$ over the entire region, an order of magnitude better than previous maps. At each position on the map we use the median colour-indices of the stars within a $10''$ box to calculate the local extinction law parameter α using the equation:

$$\frac{\langle \lambda_1 - \lambda_2 \rangle}{\langle \lambda_2 - \lambda_3 \rangle} = \frac{\left(\frac{\lambda_2}{\lambda_1} \right)^\alpha - 1}{1 - \left(\frac{\lambda_2}{\lambda_3} \right)^\alpha}$$

Until recently, this extinction law parameter which indicates the relationship between extinction and wavelength as $A_\lambda \propto \lambda^{-\alpha}$ was thought to be a constant $\alpha \sim 2$ (Martin & Whittet




Figure 1. Histograms of the values of extinction law parameter α measured across the NB (*left* figure) and the absolute extinction in magnitudes in the J , H and K -bands (*blue*, *green* and *red* respectively in the *right* figure). The fit to the measured α is shown as a gaussian with mean of 2.20 and standard deviation of 0.18.

1990; Rieke & Lebofsky 1985; Nishiyama et al. 2006) for the NB. We show that it is in fact highly variable in the NB. Using a local extinction law calculation, we then measure the absolute extinction in the J , H , and K -bands using the equations:

$$A_{\lambda_1} = \frac{\langle \lambda_1 - \lambda_2 \rangle}{1 - \left(\frac{\lambda_1}{\lambda_2} \right)^\alpha} \quad A_{\lambda_2} = \frac{\langle \lambda_1 - \lambda_2 \rangle}{\left(\frac{\lambda_2}{\lambda_1} \right)^\alpha - 1}$$

Our final calculation of absolute extinction takes into account the variation in the extinction law as well as variation in the amount of absolute extinction at each location. Using data from the 3 bands, J , H , and K , we are able to calculate three measures of the extinction law and two measures of the absolute extinction for each band at each position in our map. Full details of this method will be presented in Gosling et al. (2008).

3. Measured extinction

Figure 1 shows histograms of the value of the extinction law parameter α and the values of absolute extinction measured from the three colour-indices ratios.

The same distribution of α is measured from all three colour indices ratios, and this distribution can be described by a gaussian with mean 2.20 and standard deviation 0.18. We suggest that a specific extinction law calculation be undertaken for all observations in the NB if possible, however, when this is not possible, we recommend a value of $\alpha = 2.2 \pm 0.54$ be used. In such cases the possible variation from this mean of α should be considered as it can have a large effect on the relative values of absolute extinction calculated.

Using the specific, local extinction law, we calculated the value of absolute extinction in the J , H and K -bands from all three colour indices ratios. We measured the extinction in magnitudes to be $1.8 < A_J < 10.0$, $0.7 < A_H < 7.0$ and $0.3 < A_K < 6.0$. All three distributions have high extinction tails extending to higher values of absolute extinction than these ranges. Figure 2 shows the resultant extinction map as well as a false colour NIR and mid-infrared image of the same region for comparison.

We repeated the extinction calculation described above using a single extinction law. We used a χ^2 -test comparison between the extinction calculated for different bands to show that the calculation using the variable extinction law produced statistically consistent extinction in all three bands (differences within 3σ of the overall distribution), whereas using a single extinction value produced statistically inconsistent extinction between bands (differences of 10 - 50σ of the overall distribution).




Figure 2. *Main Figure:* A_K map of the Nuclear Bulge made using data from the UKIDSS-GPS survey (see text) with resolution of $5''$. The colour scale ranges from 0 (white) to 3.5 (black) magnitudes of extinction. Square black regions are area for which there is no data as yet in the UKIDSS catalogue. There is turnover to low extinction within some regions of high extinction where only foreground stars are observed as all background NB stars are obscured. *Bottom-left:* J , H , and K -band false-colour image of the NB from the UKIDSS-GPS, the photometry of which was used to produce the extinction map. *Top-right:* *Spitzer* false-colour image of the same region using the 3.6, 5.4 and $8.0\ \mu\text{m}$ channels scaled to reveal dust emission. Note the correspondence between the emission in this image and the regions of high extinction from the UKIDSS map.

4. Conclusions

We have presented a new, highly detailed map of the NIR extinction towards the NB using data from the UKIDSS-GPS. We find both variation in the degree of absolute extinction, and in the extinction law parameter α . We show that the extinction law is not “universal” as had previously been thought (Martin & Whittet 1990; Rieke & Lebofsky 1985), but is highly variable, and that only when this variation is taken into account can absolute extinction be calculated consistently at different wavelengths. We recommend that this variable extinction law be taken into account in all future extinction corrections. We intend to compare our extinction law and absolute extinction maps to dust emission observed with *Spitzer* (*top-right* insert in Figure 2) to better determine the properties of the extinguishing material and so the distribution of gas and dust in the Bulge.

References

- Catchpole, R. M., Whitelock, P. A., & Glass, I. S. 1990, MNRAS, 247, 479
- Dutra, C. M., Santiago, B. X., Bica, E. L. D., & Barbuy, B. 2003, MNRAS, 338, 253
- Lucas, P. W., et al. 2007, ArXiv e-prints, 712, arXiv:0712.0100
- Rieke, G. H., & Lebofsky, M. J. 1985, ApJ, 288, 618
- Martin, P. G., & Whittet, D. C. B. 1990, ApJ, 357, 113
- Nishiyama, S., et al. 2006, ApJ, 638, 839
- Schultheis, M., et al. 1999, A&A, 349, L69